

GREENHOUSE GAS EMISSIONS BY AREA OF SYDNEY

REGION: Area	Tonnes per Capita Annually	Compared to National Average
CENTRAL	26.85	42.1%
Leichhardt	27.03	43.0%
Botany Bay	19.73	4.4%
Sydney-Inner	37.11	96.3%
Marrickville	23.54	24.6%
EAST	25.76	36.3%
Waverly	25.35	34.1%
Randwick	23.36	23.6%
Woollahra	28.56	51.1%
NORTH	22.20	17.5%
Hornsby	22.07	16.8%
Baulkham Hills	20.54	8.7%
Manly	24.02	27.1%
Ku-rin-gai	25.32	34.0%
Pittwater	20.55	8.7%
Warringah	20.70	9.5%
INNER NORTH	26.48	40.1%
North Sydney	30.62	62.0%
Mosman	27.75	46.8%
Ryde	22.29	17.9%
Willoughby	25.97	37.4%
Lane Cove	25.75	36.2%
SOUTH	19.69	4.2%
Sutherland Shire West	18.85	-0.3%
Rockdale	19.72	4.3%
Sutherland Shire East	19.76	4.6%
Kogarah	20.41	8.0%
SOUTHWEST	17.19	-9.0%
Campbelltown	16.68	-11.7%
Camden	17.69	-6.4%
WEST	18.11	-4.2%
Liverpool	17.63	-6.7%
Auburn	19.38	2.5%
Holroyd	18.39	-2.7%

Fairfield	17.47	-7.6%
Bankstown	17.42	-7.8%
Parramatta	19.88	5.2%
Penrith	17.10	-9.5%
Blacktown N	18.30	-3.2%
Blacktown SE	17.92	-5.2%
Blacktown SW	17.61	-6.8%
INNER WEST	21.65	14.6%
Drummoyne	22.75	20.4%
Hurstville	19.53	3.3%
Canterbury	19.04	0.7%
Concord	21.22	12.3%
Strathfield	23.12	22.3%
Burwood	22.08	16.8%
Ashfield	23.83	26.1%
NEW SOUTH WALES AVERAGE	19.31	2.2%
NATIONAL AVERAGE	18.90	0.0%

From the Australian Conservation Federation "Consumption Atlas"
http://www.acfonline.org.au/custom_atlas/index.html

Regional averages are unweighted (not weighted by population)

www.demographia.com
20070712